W.345

TL 568 .A21a #1/80

# A PRELIMINARY STUDY OF UNIDENTIFIED TARGETS OBSERVED ON AIR TRAFFIC CONTROL RADARS

By

Richard C. Borden, Electronics Division and

Tirey K. Vickers, Navigation Aids Evaluation Division Technical Development Report No. 180



CIVIL AERONAUTICS ADMINISTRATION
TECHNICAL DEVELOPMENT AND
EVALUATION CENTER
INDIANAPOLIS, INDIANA

May 1953

FAA Library

## U. S. DEPARTMENT OF COMMERCE Sinclair Weeks, Secretary\_

## CIVIL AERONAUTICS ADMINISTRATION

F. M. Lee, Administrator

D. M. Stuart, Director, Technical Development and Evaluation Center

FEDERAL AVIATION AGENCY Class ~ Book . AZIN Vol Copy 3

This is a technical information report and does not necessarily represent CAA policy in all respects.

#### TABLE OF CONTENTS

	 Page
SUMMARY	
INTRODUCTION	
OFFICIAL RECORDS	
HISTORICAL REFERENCES	
WASHINGTON OBSERVATIONS	
ANALYSIS OF WASHINGTON DATA	
SUPPLEMENTARY OBSERVATIONS AT INDIANAPOLIS	
ANALYSIS OF SUPPLEMENTARY OBSERVATIONS	
EFFECT ON AIR TRAFF IC CONTROL OPERATIONS	
CONCLUSIONS	 14
RECOMMENDATIONS	 15
BIBLIOGRAPHY	 16

The Air Navigation Development Board (ANDB) was established by the Departments of Defense and Commerce in 1948 to carry out a unified development program aimed at meeting the stated operational requirements of the common military/civil air navigation and traffic control system. This project, sponsored and financed by the ANDB is a part of that program. The ANDB is located within the administrative framework of the Civil Aeronautics Administration for housekeeping purposes only. Persons desiring to communicate with ANDB should address the Executive Secretary, Air Navigation Development Board, Civil Aeronautics Administration, W-9, Washington 25, D. C.

#### A PRELIMINARY STUDY OF UNIDENTIFIED TARGETS OBSERVED ON AIR TRAFFIC CONTROL RADARS

#### SUMMARY

This report describes the investigation of a type of unidentified moving target which has been observed recently in considerable numbers on the viewing screens of air traffic control radar equipment operated by the Civil Aeronautics Administration. This investigation was conducted by means of interviews with personnel concerned, by study and correlation of official records, and by first-hand observation of numerous targets on the Washington Microwave-Early-Warning (MEW) radar and on the Indianapolis ASR-2 radar.

It was determined that targets which are known to operating personnel by various terminologies such as "ghosts," "angels," or "pixies" do not represent new phenomena; nor are they peculiar to the Washington area. Correlation of controllers' reports with United States Weather Bureau records indicated that a surface temperature inversion was almost always noted when such targets appeared on the radar.

Firsthand observation in the tracking and subsequent motion analysis of 80 of these unidentified targets indicated that a large number of these were actually secondary reflections of the radar beam. Apparently these reflections were produced by isolated refracting areas which traveled with the wind at or near the temperature inversion levels.

Although the exact size, shape, and composition of these isolated areas are not known, it is believed that they may be atmospheric eddies produced by a shearing action of dissimilar air strata. It appears possible that such eddies may refract and focus the radar energy with a lens effect to produce small concentrations of ground return with sufficient intensity to show up on the radar display. It is also believed that the correlation of the appearance of these radar targets with visual reports of so-called "flying saucers" is due to the strong probability that both effects are caused primarily by abrupt temperature inversions.

Such radar targets are usually easy to recognize because of their generally weak return and slow ground speed. Unfortunately, radar returns from small helicopters sometimes present these same characteristics. Spurious targets of this type can become a nuisance under busy traffic conditions, particularly in localities where helicopter operations are prevalent.

ţ

#### INTRODUCTION

Closely related to a recent flood of visual reports of flying saucers, the sighting of scores of unidentified targets on the Washington Air Route Traffic Control Center (ARTC) radar aroused much publicity and speculation regarding the origin, composition, and import of these objects. Concerned with the possible detrimental effects of this situation on the control of air traffic, the Air Navigation Development Board requested the Technical Development and Evaluation Center of the CAA to investigate the problem.

The specific objectives of this study were:

- 1. To find out as much as possible about the nature of the targets themselves.
- 2. To determine whether this problem is new and peculiar to the Washington area or whether it had occurred previously at Washington and at other CAA radar locations.
- 3. To determine the effect of this problem on the control of air traffic.
- 4. To determine what changes should be made in the radar development program in order to cope with the situation.

#### OFFICIAL RECORDS

As one of the first steps in this study, all records of these phenomena reported in the logs of the Washington ARTC Center were tabulated. The tabulation, given as Table I of this report, was taken to the Analysis Section of the United States Weather Bureau where it was correlated with meteorological data for the periods involved. It was then discovered that a temperature inversion had been indicated in almost every instance when the unidentified radar targets or visual objects had been reported. Weather analysts were asked whether any unusual weather conditions had prevailed over the Washington area during the period covering the occurrences of large numbers of the unidentified radar targets. Their report may be condensed as follows:

Monthly Weather Summary, July 1952.

The heat wave that broke records in the eastern portion of the United States during the month of June continued on through July, becoming intensified during the latter part of the month. July weather maps were characterized by a well-developed Bermuda high

1		T	T		£	T	-		1			7-		-			<del></del>									-	
	Remarks	Speed 20 to 35 miles per hear, Followed curved course from 13 miles seems of Arcels grey Hanness, in Proc.	McLean, No details available.				Fair to weak langula, appeal 100 to 130 miles per hean	Lights proced rapidly up, down, and herisantally. Also	To details assisted		Control of the second s	Light and frieff wither opposited to faction attents from platesty of tornion the state of DCA attents.	Moretimes to bell medibant at 3 to 40 miles per bear.	strong, or the same threes, Mostly weak, occasionally attorn.	Conter reduct other largets at £200 EST,	Small etretar object, whip accasamally visible. He moise, opening at 4 to 90 to 60 miles by her heart (Destinates marries marries at Count morting and parties are naving and the second morting and the second morting and the second	Drilliant light, trempliken ispania, is	Vicinity thandoratern. Darting grand adam 1 ct.	11421	Leve, unstangy light, moving north to south. Left no trail.	Maved slowly, slopped, filchared, moved is arc.	of nature to narch-seribant heading from 6 miles seeks-seekbees	Many rightings.	Mavement from Maredon to a con-	15 miles mide.	No details available.	Round white objects,
A Redigered Observation	Humidity	Nermal	Not available for horality	DCA: Low, below measuring	limits at 11,000 ft,	Norfolk: High, but fell off at 6000 ft. DCA: Sharp fall at 6000 ft.	DCA: Above 10,000 ft, dropped, then increased sightly, dropping again at 15,000 to	DCA: Above 10,000 ft. dropped, then increased slightly, dropped,	DCA: Abere 10,000 ft. dropped.	Again at 15,000 ft. DCA: Above 10,000 ft. drugged.	again at 15,000 ft.	DCA: Above 10,000 ft. dropped, then increased slightly, dropping again at 15,000 ft.	DCA: Sharp derrease et 10,000 ft.	DCA: Fell below measuring	Umits at 8009 ft.  DCA: Mish is 12 odd in 7211	18,000 ft.	DCAt Migh to 12,000 ft., fell off comowhal, sharp rise at 18,000 ft.	Not available for locality	Not available far locality	Mak to 12 may 2 con as	semembat, sharp rice at 18,000 ft.	DCA: Migh at 10,000 ft., alow (all	15,000 ft. DCA: Wigh at 10,000 ft., slew fall to 15,000 ft., fast fall at	5,000 ft. CA: High at 10,000 ft., alow fall	15,000 ft., fast fall above 15,000 ft., DCA: High to \$000 ft., above felt	at 9000 ft.	DCA: High to \$600 ft., sharp fall, then increasing to 100 per cent
	Note Note	1506 ft., 1º from 9600 to 10,000 ft., otherwise normal		DCA: Burfaco inversion 6.	Norfells	rate around 9000 ft. DCA: Serface inversion 2"	DCA: Barface inversion 3* leathermal between 8000 and 10,000 ft.	DCA: Burface inversion 3* incidential between 8000 and	DCA: Surface invarcion 3*	DGA: Surface inversion to	DCA: Burfare interest	10,000 ft.	DGA: Burface Inversion 3*	DCA: Surface inversion I*	+	7	ī	Not available for locality	Het avatlable for locality	$\top$	18,000 R. amail tweavalon at	DCA: Burface laverales 4", of temple person rate to 10,000 ft.	DCA: Surface Inversion 4", a	+	-	-+	2000 ft., inversion 900 ft. thich the 10 2500 ft.
To the same			Wational 42	Copt. Bres	į	Ferry 901	Contac	Capital 80v	USAF Personnel	Comfer	Capital 610		Comtor	Conte	7. A	$\top$	_	American Sic	Cilian	1	1	PCA Test	Cilinens	Conter	Center	$\top$	Pilot
Hally de	3		7000	11,000	30	8										T	1			3 00 0	-			Ť	Ť	Ť	
-	×2	Are	Outmile	do Miles Southwest	Vicinity		South A DW	Me rithsburg	Over ADM	PCA To Table	PCA PCA	Arre	Terminal	DCA Terminal	Miverdale	Greenbelt		1	C. yanda ra	ADA		DCA Tarminal Area	City of Washington	DCA Terminal	DCA Terminal	Miles	Southernt ADW
j		$\perp$		2 2	3	$\perp$	$\perp$	Ì	Ore						4 8	T	T			1	Ť		0,5	1000	1	A Pile	
Tornet DEA BEA	L	_ ։	•	•	ŀ	$\perp$			•							1.	1.	•	•		†	1	*	<del>                                     </del>	T	-	+
¥   	_	+			$\vdash$	+	+	$\dashv$				I	$\Box$			I	士				$\dagger$	$\dashv$		$\vdash$	$\vdash$	+	$\dashv$
1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	-	+	-	·	$\vdash$	╬	+	$\dashv$			-	╀.	-+		<u> </u>	$\perp$	4	$\dashv$			Ŀ	$\Box$				上	
1	Fail.					T	$\dagger$	+				╁	+	-	-	+	+	$\dashv$		-	+	$\dashv$		•	-	+	4
15	1	-	_	_	-	╀	+	+	- 7	•	_	2	+	<u>.</u>	-	-	ŀ	4	-	<u> -</u>	1.			1	i	-	
25.	5-23 2000		_	-13	1-14 2112	7-19	7-20 0180			7-20 83 88 83 88	8	3			97	802	I		3	8077	1		ē	3 2 2	3 8 8 8 8	1500	
					•	17	17	1 7		~ I	7	13	7	. 1	7-27	7.17	1	- T	17.	7-27	17	_		5.	67-1	7	

TABLE ! (Centimed)

	l from alsport	·			ed of 30 miles a west of DCA.	1,1			ADV - Andress Air Posse	Base APC - Approach Centrel	CTR · Center DCA · Vashington	EFF - Eathorn standard time est Eathmaded	Mil Mean son here! TWR - Tower
	2	25,00		l	**	39	1		1	· ₹	<u> </u>	<u> </u>	
Remarks	Oblang light, Note: may have been light from airport ceiferneter.		Marring continued.	No details available.	Mering sast to southeast at average speed of 30 miles per hour. I list appeared 20 to 25 miles west of DCA. Winds to 20,000 ft, averaged 18 to 20 houts.	Class 4 targets, speed 60 miles per herr, tracked from 18 miles serth of DCA to 3 miles serth of DCA.	Heading sast.	Moring in are high everhead.	Targeta plotted on combenst and neeth bestings at 24 to 59 knots. Most largets within 10 miles of rader sevenes.	Target plotted on assi-southerst beading, speed 53 knots, curved path.	Slow-moving target.	Targets picted on north to east- northeast headings, speed 26 to 45 knots.	Targets plotted on west-northwest to north-northwest hadings, speed 21 to 43 knots.
Observations Humidity	Not available	Not available for locality	DCA: Decreasing to very dry at 14,000 ft.	DCAL High throughout	DCA: High throughout	DCA: High throughout	DCA: High, decreasing to below measuring limits at 17,000 ft.	DCA: High at surface, low above upper inversion, otherwise below limits	DCA: High at surface, for above upper inversion, otherwise below limits	DCA: High, decreasing sharply at 14,000 ft.	DCA: High, decreasing shorply at 14,000 ft.	DCA: Kigh, with abarp fluctuations between 16,000 and 21,000 fl.	DCA: High, with sharp fluctuations between 16,000 and 23,000 ft.
Redisered Observations Temperature Lapse Rate	Not available	Not available far locality	DCA: 8mail sarfere laveration isothermal at 11,860 ft., small inversion at 14,800 ft.	DCA: Small surface inversion	DCA: Small surface inversion	DCA: Normal	DCA: Normal	DCA: Surface treesides Julium 2000 fly, another between 3000 and 1000 fly.	DCA: Sector inversion below 2006 ft., another between 2000 and 9800 ft.	DCA: Surface inversion 6", upper inversions at 13,300 and 15,000 ft.	DCA: Surface inversion 6', upper inversions at 13,900 and 15,000 ft.	DCA: Sarface inversion to	DCA: Surface inversion to 408 ft., isothermal to 1108 ft.
Reported By	Lecal	Local	Capital 982	ADW Approach Control	Center	DCA Tower	Crater	Lecal	Contor	Contor	ADW	Conter	Conter
Altitude	(Jeef)		19,000										
Location	City of Washington	25 Miles Nacib Savege, Md.	South South DCA	DCA Terminal Area	DCA Termismi Aree	DCA Terminal Ares	DCA Terminal Area	City of Weshington	DCA Terminal Area	15 Miles West DCA	1 1/2 Miles Seedbesel ADE	DCA Terminal	DCA Terminel Area
300			316	,				a Mir					
Vismi Combel	•	•	•										_
TO	NA A	T						<u>  .                                     </u>		_		_	
Bed Scales	<u> </u>					•					1_	1_	-
1	CTR			_	-			-	<u> •</u>	<u> •</u>	-	<u>  •                                     </u>	<u> •</u>
Number Targets	_	_	_	1	i	_	_	-	:	<u> -</u>	-		-
1	• • • • • • • • • • • • • • • • • • • •	i	1	3 0000	3 00 0	8.	:	8 -7	3 000	***	5693	1	3 2 2 2
200	2	1.5	2	1	1	1	2	3	3	i	7.	1	ä
L													

pressure area which remained in the vicinity of the southeastern coast line during the entire period. This high pressure area was responsible for an anticyclonic (clockwise) circulation of air over the eastern United States, a movement which continued during the month. This flow brought warm, moist air up from the Gulf of Mexico. The warm air mass usually extended up to about 10,000 feet. At higher levels the flow was from the west-southwest, and this continental air mass from the southwestern desert and drought area was hot and dry. Stagnation and heating of the air over the eastern United States was further increased because of an extremely strong band of westerly winds along the northern United States border, winds which prevented cold Canadian air masses from pushing south. Cyclonic activity was confined mostly to the area north of this band of westerly winds. There was a notable lack of thunderstorm activity in the Washington area. Physicists at the Naval Observatory reported that the amount of electrification in the air was very low.

The foregoing analysis indicated that the lack of cloud cover promoted solar heating in the daytime and rapid radiation cooling of the surface at night. This combination, with the prevailing light winds, was unusually conducive to the formation of temperature inversions during the hours of darkness.

Since the visual reports of flying saucers indicated that the observed lights spanned the same color range as the aurora borealis and since auroral effects closely follow sunspot activity, personnel of the Naval Observatory were consulted in order to determine whether any unusual sunspot activity had occurred during the period in question. They reported that there had been no unusual activity of this nature.

Reports from Other Locations.

The Washington ARTC Center is the only one equipped with air route surveillance radar. However, several CAA control towers are equipped with airport surveillance radar, Type ASR-1. A survey of these locations produced the following results:

- ATLANTA, Municipal Airport. No unidentified targets of this nature have been reported.
- BOSTON, Logan Field. Unidentified targets have been noticed on rare occasions. One slow-moving target was observed during instrument flying weather conditions about August 1, 1952. No interference with traffic has been caused by this problem.

- CHICAGO, Midway Airport. Unidentified targets have been seen on many occasions, particularly when temperature inversions have been in effect and low smoke hung over the city. They are usually given as traffic information to other aircraft and occasionally form a nuisance problem, since there is a considerable helicopter activity at and around the airport.
- CLEVELAND, Municipal Airport. Unidentified radar targets have been observed many times. The chief controller reported that on a recent occasion such targets moving slowly from west to east showed up in all portions of the scope face.
- MINNEAPOLIS, International Airport. No targets of this nature have been reported.
- NEW YORK, New York International Airport.

  No targets of this nature have been reported.
  - La Guardia Airport. Only one such instance was reported. At the time it was thought to be due to difficulties within the radar itself.
- WASHINGTON, National Airport. Targets of this nature have been observed occasionally over a long period. Recent occasions are logged in Table I of this report.

#### HISTORICAL REFERENCES

The history of radar abounds with reports of strange echoes received from supposedly clear skies. Early observers suspected birds or stray weather balloons, but these were eliminated by visual checks. Conjecture that clouds of insects were responsible was also eliminated when such echoes were obtained in the dead of winter. Some connection with the weather was suspected after it was noted that echoes of this type became more numerous on summer nights under calm conditions. Additional evidence indicated that many of these echoes originated in the fine structures of the dielectric (refracting) layers of air-mass boundaries and in regions of air turbulence. Some of the sharpest echoes involved surfaces of pronounced transitions of the water-vapor content of the air. The bibliography at the end of this report contains numerous detailed references to these general phenomena.




Fig. 1 Track Plots of Unidentified Targets, Washington MEW Radar, 1958 to 2242 EST, August 13, 1952

WASHINGTON OBSERVATIONS

August 13-14, 1952.

The observation period started at 1830 Eastern standard time (EST) on the evening of August 13. At the beginning of this period, the moving target indicator was gated to cancel out ground returns up to a range of 10 nautical miles. Beyond this range the scope was clear except for a few permanent echoes that were visible.

Suddenly, at approximately 1957 EST, a group of seven strong stationary targets became visible in an area about 15 miles north-northeast of the radar antenna. During the next two or three antenna revolutions, the area on the scope between Washington and Baltimore became heavily sprinkled with stationary targets in a belt about 6 miles wide. A group of additional targets became visible in an area approximately 10 to 15 miles south of the radar antenna. This was evidence of the beginning of a temperature inversion.

Within the next minute, at approximately 1958 EST, four unidentified moving targets showed up 5 miles southeast of the radar antenna and moved in a southerly direction away from it. When the radar beam was

Fig. 2 Track Plots of Unidentified Targets, Washington MEW Radar, 2242 to 2352 EST, August 13, 1952

switched from high to low, the targets disappeared. The beam was switched back to high, and the targets returned.

Targets were uniformly small and usually had a weak, fuzzy appearance. However, the target intensity varied from sweep to sweep. Occasionally one or two very strong returns would be received in succession, followed by almost total blanking.

For the next four and one-half hours, many unidentified targets were carefully plotted with a grease pencil on the face of the Type VG scope. The time for each was entered on these plots in order to calculate ground speeds. To secure a permanent record, time data and track plots were transferred from the scope face to a sheet of frosted acetate. These plots are reproduced in Figs. 1, 2, and 3. The distribution of target ranges is shown in Fig. 4. The average distance that any target was tracked continuously was approximately 2.1 nautical miles.

The observation period was discontinued at 0030 EST on August 14, and steps were taken to secure all available meteorological data relevant to the observation period. The local radiosonde observation




Fig. 3 Track Plots of Unidentified Targets, Washington MEW Radar, 2353 to 0029 EST, August 13-14, 1952

which was taken near the midpoint of the observation period, at 2200 EST on August 13. is reproduced in Fig. 5. Winds aloft, as observed at the same time, are listed in Table IL

August 15-16, 1952.

On the night of August 15-16, additional track plots were obtained by Washington ARTC Center personnel. During this period, the radar was operating on the high beam with the moving target indicator gated to 12 miles. The same stationary targets in the Washington-Baltimore belt and in an area 10 to 15 miles south of the radar antenna were visible again on the scope face.

Track plots for this period are shown in Figs. 6 and 7. The local radiosonde observation taken at 2200 EST on August 15 is reproduced in Fig. 8. Winds aloft, as observed at the same time, are listed in Table III.

#### ANALYSIS OF WASHINGTON DATA

It will be noted from Table I that many more unidentified targets are picked up by the Washington ARTC Center than by the Washington Airport Traffic Control Tower. This may be explained by the fact that the center is equipped with a MEW radar, while the tower is equipped with an airport surveillance radar, Type ASR-1. The most significant differences between the two types of equipment are listed in the following:

1. The peak power of the MEW is 3

decibels (db) higher than the ASR-1.

2. The average power of the MEW is 6 db higher than the average power of the ASR-1.

3. The MEW has a higher elevation angle

coverage.

4. The MEW elicits approximately twice as many hits per scan per target since the scan rate of the MEW is 6 revolutions per




Fig. 4 Distribution of Target Ranges, Washington MEW Radar, August 13-14, 1952 Observation

minute (rpm). Additional specifications of these radars are listed in Table IV.

The almost simultaneous appearance of the first moving targets with the ground returns, signifying the beginning of the temperature inversion, suggested that the target display was perhaps caused by some effects existing in or near the inversion layers.

It will be noted in Figs. 1, 2, and 3 that all targets observed in the first period were moving from the north or northwest. In Fig. 6 all targets were moving from the south or southwest, and in Fig. 7 all were moving from the west or northwest. The definite directional trend in each case eliminated the possibility that the unidentified targets were

ity wide t	are undersease			Surface	170	5
				1000	180	24
	MADLEII			2000	190	26
	TABLE II		•	3000	210	24
721	INDS ALOF	·T		4000	210	23
W A CTTTN	GTON (SILV	rn unil		5000	220	20
		t 13, 1952		6000	220	16
2200 EST	. Augus	1 13, 1752		7000	220	18
A 14244	Dimension	Velocity		8000	220	17
Altitude	Direction	•		9000	220	13
(MSL)	(Degrees)	(Knots)		10000	240	12
•	. •			11000	270	11
Surface	Calm	0		12000	270	13
1000	Calm	0		13000	260	17
2000	350	12		14000	260	21
3000	340	12		15000	260	25
4000	320	14		16000	270	25
5000	320	16		17000	270	23
6000	300	18		18000	270	22
7000	300	20	•	19000	270	21
8000	310	20	• 1	20000	260	20
9000	310	22		21000	270	22
10000	300	26		22000	280	24
11000	290	28		23000	290	26
12.000	290	29		24000	280	26
13000	300	30		25000	290	26
14000	300	28		26000	300	30
15000	290	29		27000	300	34
16000	300	29	<u>-</u>	28000	300	38
17000	300	29		290 <b>00</b>	290	38
18000	300	30		30000	29 <b>0</b>	36
19000	300	32		31000	300	35
20000	300	38		32000	300	35
21000	290	38		33000	310	34
22000	280	43		34000	310	40
23000	280	48		35000	300	47
24000	280	50		36000	300	49
25000	270	52		37000	300	50
26000	280	57		38000 -	300	48
27000	270	61		39000	310	42
28000	270	54		40000	320	38
29000	270	55		41000	300	43
30000	280	62		42000	300	53
31.000	270	63		43000	300	67
32000	280	73		44000	310	- 69
33000	280	84		45000	310	<i>.</i> 60
-						

surface vehicles such as trains, trucks, automobiles, or boats. Had this been the case, some vehicles would have been moving in the reverse directions. In each case, target directions corresponded with the wind

TABLE III

WINDS	ALOFT
WASHINGTON	(SILVER HILL)
2200 EST	August 15, 1952

Altitude Direction Velocity

(Degrees)

(Knots)

(MSL)




Fig. 5 Radiosonde Observation, Silver Hill, Washington, D. C., 2300 EST, August 13, 1952




Fig. 7 Track Plots of Unidentified Targets, Washington MEW Radar, 2253 to 0450 EST, August 15-16, 1952

directions reported aloft. This fact suggested that whatever was producing the targets was being carried by the wind.

The next step of the analysis was to determine, if possible, the altitude of the objects which produced the radar targets.




Fig. 6 Track Plots of Unidentified Targets, Washington MEW Radar, 2213 to 2244 EST, August 15, 1952




Fig. 8 Radiosonde Observation, Silver Hill, Washington, D. C., 2200 EST, August 15, 1952

Since the radar actually measures slant range which could in some cases be almost directly overhead from the high-beam MEW antenna, the minimum range of each target was used to determine the absolute maximum altitude of the object producing the target.

#### RADAR EQUIPMENT CHARACTERISTICS

	Tower Radar	Center Radar
Type	ASR-1	MEW
Frequency	S-band	S-band
Pulse-repetition frequency	1,000	900
Pulse rate	0.5 microsecond	l microsecond
Vertical coverage	6,000 feet at 6 miles	12,000 feet at 3 miles
Scan Rate	28 per minute	6 per minute
Display scopes	12DP7	12DP7 and VG2
Power output	200 kilowatts	400 kilowatts

For example, a target which came within five nautical miles of the radar antenna could not be above an altitude of five nautical miles, or 30,400 feet. With the use of the slant-range principle, the absolute maximum altitude of each target was determined and is listed in Table V. When attempting later to determine the probable altitude of each target by studying the winds aloft, it was useful to have these maximum altitude figures to eliminate the necessity for consideration of higher altitude levels.

Since winds aloft can vary considerably during the period of a few hours, it was decided to use in this analysis only data on targets which were under observation during the periods from one hour before to one hour after the observations of the local winds aloft. These targets are listed in Table V.

During the observation period on the night of August 13-14, all targets on a southerly heading had ground speeds of at least 24 knots. The only reported winds with a southerly heading had a velocity of only 12 knots. These were winds at the 2,000- and 3,000-foot levels. Targets on a southeasterly heading had a speed range of 32 to 48 knots. However, the only winds on this heading were from 14 knots at 4,000 feet to 38 knots at 20,000 feet.

During the August 15-16 observations, targets on a north or northeasterly heading had speeds of 35 to 42 knots. The only reported winds moving in this direction ranged between 5 and 26 knots from the surface up

to 9,000 feet. Targets on easterly headings had speeds from 22 to 45 knots. The only reported winds moving in this direction had speeds of from 10 to 24 knots between 10,000 and 25,000 feet.

In Figs. 9 and 10, the directions and velocities of the winds aloft are plotted on a polar projection diagram together with the directions and velocities of the observed targets. Agreement between the directions of the winds and the directions of the targets is apparent.

One of the theoretically possible causes of the unidentified targets was the delayedpulse or second-time-around effect inherent in the radar method of time measurement. With a second-time-around effect, objects beyond the normal sweep range of a radar can be displayed on the scope because of reception of an echo pulse elicited not by the transmitted pulse which triggers the range sweep but by the preceding transmitted pulse. The apparent velocity of the target on the radar is no greater than and normally less than the velocity of the object producing the return. The heading of the radar target would not necessarily be parallel to the heading of the object unless the object was on a course radial to the radar antenna. These effects are illustrated in Fig. 11.

If we assume then that an object producing a second-time-around radar target was being carried by the wind, the apparent velocity of the target would be no greater than the wind velocity. However, the analysis of the targets listed in Table V showed that

MOVEMENT DATA ON TARGETS TRACKED WITHIN ONE HOUR FROM START OF OBSERVATIONS OF WINDS ALOFT

Date Aug. 1952	Starting Time EST	Direction (Degrees)	Target Speed (Knots)	Reflector Speed (1/2 Target Speed)	Absolute Maximum Altitude (Based on Minimum Slant Range	Probable Altitu (Based on Winds Aloft)
13	2159	005	28	14	63000	2000
	2201	360	24	12	75000	2000
	2229	310	33	16.5	23000	8000
	2240	300	46	23	30000	9000
	2242	325	48	2 <b>4</b>	33000	9000
	2259	010	31	15.5	31000	2000
	2303	330	42	21	36000	8000
	2330	340	39	19.5	23000	5000
	2330	305	39	19.5	24000	8000
	2331	315	39	19.5	35000	8000
	2332	315	36	18	23000	8000
	2345	310	38	19	19000	8000
	2347	310	42	21	43000	8000
	2349	290	39	19.5	35000	7000
	2356	300	42	21	37000	7000
	2355	350	36	18	83000	2000
15	2213	260	45	22.5	34000	14000
	2226	225	35	17.5	24000	900
	2230	250	28	14	37000	10500
	2238	185	36	18	29000	900
	2240	210	42	21	18000	4500
	2353	275	23	11.5	29000	10500*

<sup>\*</sup>This target could also have been a direct radar return from an object floating with the wind a 15000 to 17000 feet mean sea level.

they were actually moving at speeds approximately double the wind velocities reported for the directions involved. This fact eliminated the possibility that the targets were being produced by the second-time-around effect.

When the target velocities plotted in Figs. 9 and 10 were halved, those plotted points clustered very closely around the wind plots. Further investigation of the doubledspeed effect indicated that this effect could be produced if the original radar beam were reflected downward to give a ground return, as shown in Fig. 12. If we assume that some sort of horizontal reflector was present aloft and that the angle of reflection equalled the angle of incidence of the radar beam, any horizontal movement of the reflector would produce a movement twice as great in the image being received on the radar scope. Furthermore, the apparent motion of the image would be parallel to the motion of the reflector, as illustrated in Fig. 13.

When the observed target velocities were divided by two, the target motions

corresponded closely to the reported win directions and velocities at certain altitud levels. In nearly all of these cases th altitude levels, which are listed as probabl altitudes in Table V, were at or adjacent the temperature inversion levels.

With only one exception, no target were seen moving at the speed and headin of the reported wind at any altitude. Thi suggested that the reflecting areas, whic were capable of bending the radar beam, wer nevertheless not of sufficient density t produce direct returns on the radar scope Thus, it appeared likely that the reflectio effect was being produced by the atmospher itself. If this were the case, it woul probably be a refraction rather than reflection which was involved. This effect i shown in Fig. 14.

The uniformly small size of the observed targets as well as the relativel low frequency of their occurrences suggester that the conditions producing this effect were extremely localized and decidedly critical Although the exact nature of the discontinuit




Fig. 9 Comparison Between Winds Aloft and Target Data, August 13, 1952 Observation

is not known, one possible explanation might be that it is an eddy in the atmosphere. Such eddies may be produced by the shearing effect of dissimilar air masses moving at different speeds and headings at or near the inversion boundary. They might under certain conditions produce bulges in the inversion layer, concentrating and directing the radar energy with a lens effect to produce a return signal strong enough to show up on the radar scope. The relatively short paths of some of the radar targets before their

fade-out might be attributed to the dissipation of these eddies in the stratified air mass.

Intermediate speed checks on numerous targets indicated that individual velocities remained quite steady during the observation period. It became possible to predict with accuracy the progress of specific targets from minute to minute. There was no evidence of hovering or of sudden increases in speed by any target. It is believed that previous reports of sudden accelerations of targets to supersonic velocities were due to




Fig. 10 Comparison Between Winds Aloft and Target Data, August 15, 1952 Observation

a controller's transfer of identity from a faded target to another target which was just appearing on a different section of the scope.

It would be unwise to assume that all unidentified slow-moving radar targets are caused by refraction of radar energy. Small rain clouds produce much the same appearance on the scope. Other targets could be direct returns from bird formations, balloons, or debris carried aloft by convection or tornadoes. It has recently been reported that

more than 4,000 balloons are released in the United States every day by Government and civilian research organizations. A recent analysis of more than 1,000 visual reports of unidentified flying objects by the Air Technical Intelligence Center at Wright-Patterson Air Force Base indicates that

1"Many Potential 'Saucers,'" Science News Letter, Vol. 62, No. 7, Aug. 16, 1952, p. 106.




Fig. 11 Second-Time-Around Effect

21.3 per cent of these may be attributed to balloons.<sup>2</sup>

Examination of the logs of the Washington ARTC Center indicates that there is considerable correlation between the appearance of unidentified targets on the radar scope and the receipt of numerous visual reports of flying saucers. It should be noted that abrupt temperature inversions aloft can refract light in much the same way as radar waves and produce mirage effects. In a standard reference work on meteorology,3 Humphreys reports that a temperature inversion (near the surface) of 1°C per meter bends down a light ray into an arc whose radius is 0.16 that of the earth; an inversion of 10°C per meter gives an arc radius of 0.016 that of the earth, or approximately 60 miles. This effect makes it possible for an observer to see in the sky the sun or some other bright light that is actually well below the observer's horizon. On rare occasions, multiple images of the same object may be visible. It is believed that many visual sightings of flying saucers can be explained by this phenomenon.

<sup>2</sup>"Unidentified Aerial Objects Receive Careful Analysis by Air Force Experts," The Aircraft Flash, published by Department of the Air Force, Air Defense Command, Vol. 1, No. 4, Jan. 1953, p. 4.

<sup>3</sup>Humphreys, W. J., "Physics of the Air," McGraw-Hill Publishing Company, New York, 1940.




Fig. 12 Profile View Showing Effect of Moving Reflector

## SUPPLEMENTARY OBSERVATIONS AT INDIANAPOLIS

November 4, 1952.

During test runs of the new ASR-2 radar equipment, a large number of unidentified moving targets appeared on the scope at approximately 4 p. m. The sun was low in the sky, and the sky was clear of all clouds. Ceiling and visibility were unrestricted. Pilot temperature reports from a departing aircraft indicated that a pronounced temperature inversion existed at the 6,000-foot level.

Although no targets were plotted, a check on several indicated that their movement corresponded to the direction of the wind at the inversion level, with a velocity roughly double the wind velocity. Targets were larger, stronger, and more numerous than those observed by the writers during the Washington observations. At times the clutter made it difficult to keep track of actual aircraft targets on the scope.

November 5, 1952.

At approximately 4 p. m., a group of similar targets appeared on the Indianapolis ASR-2 scope. Again the sky was clear of clouds; ceiling and visibility were unrestricted. Targets were strong, numerous, and of various shapes and sizes.

A simultaneous check of the L-band radar showed that only a few targets were being picked up by this equipment. The L-band targets appeared considerably weaker than those seen on the ASR-2 scope, although L-band aircraft targets appeared normal.

By manipulation of the ASR-2 antenna motor switch, it was possible to slew the antenna to beam it directly at some of the unidentified targets. The video return was displayed on an A-scope for closer analysis of the target characteristics. Comparisons were made with the A-scope characteristics of aircraft targets.




Fig. 13 Plan View of Reflection Effect

Aircraft targets showed sharp rise and decay times as well as relatively constant shape and amplitude. The unidentified targets showed gradual rise and decay times; amplitude and shape showed wide variations, which resulted in a random interlaced signal envelope similar to that returned by rain and cloud formations. These target characteristics are sketched in Fig. 15.

## ANALYSIS OF SUPPLEMENTARY OBSERVATIONS

The reduced target returns from the L-band radar indicated that the reflecting areas are formed by atmospheric disturbances or discontinuities rather than by some form of ionization. If the cause were ionization, it would be expected that the lower frequency of the L-band equipment would increase the susceptibility of the radar energy to reflection or refraction effects. An example of this trend is that of ionospheric layers which produce no appreciable reflection of ultra-high-frequency energy but cause strong skip propagation of the lower radio frequencies.

### EFFECT ON AIR TRAFFIC CONTROL OPERATIONS

The generally weak and fuzzy appearance as well as the slow speed of spurious radar targets usually enable them to be




Fig. 14 Refraction of Radar Beam

recognized as such by experienced radar controllers. Normally these targets have but little effect on traffic control, because they occupy very little space in relation to the entire scope area and their progress on course is very slow. The most dangerous possibility from the traffic control standpoint is the chance that one of these targets might be a helicopter.

If their course will not collide with that of an aircraft target, such targets are generally disregarded. If the course will collide with an aircraft target, some control action is indicated because of the helicopter hazard. In such cases, prudent controllers will give traffic information to pilots regarding the unidentified target, particularly at night under visual flight rule conditions. Where a collision course is involved, pilots would rather be warned about a spurious target than not be warned about a legitimate one.

At the present time, very little instrument flying is done by helicopters. Therefore, unidentified targets of this type are not usually given as traffic information to pilots known to be operating on instruments.

#### CONCLUSIONS

1. It is believed that most of the unidentified targets observed on the Washington MEW radar during the period beginning on the night of August 13, 1952 and the period beginning on the night of August 15, 1952 were ground returns caused by reflection phenomena closely connected with the temperature inversions in the lower atmosphere.

2. Unidentified radar targets of the type described in this report have been noticed since the early days of radar. Unusual weather conditions prevailing in the Washington area during the summer of 1952 were exceptionally conducive to the formation of these phenomena.

3. Present evidence indicates that the appearance of unidentified targets of this nature on radar scopes has but little effect on the control of air traffic. At its worst, it forms a nuisance by cluttering the scope

display and by requiring that additional traffic information or heading instructions be issued in order to protect other traffic against the possibility that such a target might be a helicopter.

4. In some cases, it would be desirable to provide the controller with a more positive method of identifying targets such as these so that he could determine quickly whether they are spurious or whether they are actual aircraft.

#### RECOMMENDATIONS

- l. In order to secure additional evidence regarding the causes, extent, and effects of this type of phenomena, it would be desirable to secure additional target plots from the horizontal plotting scope of the Washington ARTC Center. It would also be desirable for all CAA air traffic control agencies which use radar equipment to log the occurrence of such targets. Notes regarding the extent and motion characteristics of them, together with their effects on the control of air traffic, would also be of value. It would be desirable to correlate all these reports with official United States Weather Bureau records.
- 2. Should additional research regarding these phenomena be undertaken, close coordination with the local office of the United States Weather Bureau is essential in order that observations can be made when conducive meteorological conditions are expected.
- 3. It is believed that more complete evidence could be obtained through the use of more flexible radar equipment. A tremendous asset in evaluating the nature of false targets would be the ability to track continuously a specific target through use of a manual slewing control. It would then be desirable to examine this target closely on an A-scope radar presentation. A number of commercially available synchroscopes are ideally suited for this purpose. The echo could be enlarged on such a presentation to a width of one inch or more. Examination of the resulting trace including such characteristics as steepness of rise and decay time, energy distribution, and fluctuations in amplitude




Fig. 15 Comparative A-Scope Target Displays

should make it possible to deduce a great deal regarding the source of the reflection.

- 4. Additional simultaneous observations of the phenomena on L-band and S-band radar equipment would be desirable. The availability of aircraft which could be guided by radar to the area of the target or to the primary reflecting area would also be advantageous. Additional information may be obtained by equipping the aircraft with an aero-psychograph as well as with suitable apparatus for measuring electrical charges in these areas.
- 5. When helicopter traffic becomes more prevalent, it may be desirable to provide the controller with some type of radar accessory which can detect propeller modulation and which can give him the means to determine positively whether an unidentified target is an aircraft or a reflection. It is recommended that this type of accessory be studied in connection with the proposed evaluation program for the ASR-2 radar.

#### BIBLIOGRAPHY

- "Echoes from the Atmosphere," Bell Laboratories Record, Vol. 25, No. 2, Feb. 1947, p. 75.
- Friend, A. W., "Theory and Practice of Tropospheric Sounding by Radar," Proceedings of the Institute of Radio Engineers, Vol. 37, No. 2, Feb. 1949, p. 116.
- Friis, H. T., "Radar Reflections from the Lower Atmosphere," Proceedings of the Institute of Radio Engineers, Vol. 35, No. 5, May 1947, p. 494.
- Goldstein, Herbert, "Origin of the Echo," in "Propagation of Short Radio Waves," Massachusetts Institute of Technology Radiation Laboratory Series, McGraw-Hill Publishing Company, New York, 1951, Vol. 13 edited by Donald E. Kerr, Chapter 7, pp. 593-595.

- Gordon, W. E., "A Theory of Radar Reflections from the Lower Atmosphere," Proceedings of the Institute of Radio Engineers, Vol. 37, No. 1, Jan. 1949, p. 41.
- Gould, William B., "Radar Reflections from the Lower Atmosphere," Proceedings of the Institute of Radio Engineers, Vol. 35, No. 10, Oct. 1947, p. 1105.
- Humphreys, W. J., "Physics of the Air," McGraw-Hill Publishing Company, New York, 1940.
- "Many Potential 'Saucers,' " Science News Letter, Vol. 62, No. 7, Aug. 16, 1952, p. 106.
- "Radar Returns from the Lower Atmosphere Viewed on an AN/GPN-2 Screen," Landing Aids Experiment Station Progress Report, 1948 Test Section, Electronics, pp. 22-24.